

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317749166>

# Antiviral activity of aspalathus linearis against human influenza virus

Article in *Natural product communications* · January 2017

---

CITATIONS

5

READS

141

5 authors, including:



Ratika Rahmasari

University of Indonesia

4 PUBLICATIONS 56 CITATIONS

SEE PROFILE


Some of the authors of this publication are also working on these related projects:



Natural product activity including plant, animal and microbiota activity against human pathogen microorganism [View project](#)

# NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all  
Aspects of Natural Products Research



Volume 12. Issue 4. Pages 475-640. 2017  
ISSN 1934-578X (printed); ISSN 1555-9475 (online)  
[www.naturalproduct.us](http://www.naturalproduct.us)

**EDITOR-IN-CHIEF****DR. PAWAN K AGRAWAL**

Natural Product Inc.  
7963, Anderson Park Lane,  
Westerville, Ohio 43081, USA  
agrawal@naturalproduct.us

**EDITORS****PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry, University of Granada,  
Campus de Fuente Nueva, s/n, 18071, Granada, Spain  
afbarre@ugr.es

**PROFESSOR MAURIZIO BRUNO**

Department STEBICEF,  
University of Palermo, Viale delle Scienze,  
Parco d'Orleans II - 90128 Palermo, Italy  
maurizio.bruno@unipa.it

**PROFESSOR VLADIMIR I. KALININ**

G.B. Elyakov Pacific Institute of Bioorganic Chemistry,  
Far Eastern Branch, Russian Academy of Sciences,  
Pr. 100-letiya Vladivostoka 159, 690022,  
Vladivostok, Russian Federation  
kalininv@piboc.dvo.ru

**PROFESSOR YOSHIHIRO MIMAKI**

School of Pharmacy,  
Tokyo University of Pharmacy and Life Sciences,  
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan  
mimakiy@ps.toyaku.ac.jp

**PROFESSOR STEPHEN G. PYNE**

Department of Chemistry, University of Wollongong,  
Wollongong, New South Wales, 2522, Australia  
spyne@uow.edu.au

**PROFESSOR MANFRED G. REINECKE**

Department of Chemistry, Texas Christian University,  
Forts Worth, TX 76129, USA  
m.reinecke@tcu.edu

**PROFESSOR WILLIAM N. SETZER**

Department of Chemistry, The University of Alabama in Huntsville,  
Huntsville, AL 35809, USA  
wsetzer@chemistry.uah.edu

**PROFESSOR PING-JYUN SUNG**

National Museum of Marine Biology and Aquarium  
Checheng, Pingtung 944  
Taiwan  
pjsung@nmmba.gov.tw

**PROFESSOR YASUHIRO TEZUKA**

Faculty of Pharmaceutical Sciences, Hokuriku University,  
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan  
y-tezuka@hokuriku-u.ac.jp

**PROFESSOR DAVID E. THURSTON**

Institute of Pharmaceutical Science  
Faculty of Life Sciences & Medicine  
King's College London, Britannia House  
7 Trinity Street, London SE1 1DB, UK  
david.thurston@kcl.ac.uk

**HONORARY EDITOR****PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,  
University of Portsmouth,  
Portsmouth, PO1 2DT U.K.  
axuf64@dsl.pipex.com

**ADVISORY BOARD**

Prof. Giovanni Appendino  
Novara, Italy

Prof. Norbert Arnold  
Halle, Germany

Prof. Yoshinori Asakawa  
Tokushima, Japan

Prof. Vassaya Bankova  
Sofia, Bulgaria

Prof. Roberto G. S. Berlinck  
São Carlos, Brazil

Prof. Anna R. Bilia  
Florence, Italy

Prof. Geoffrey Cordell  
Chicago, IL, USA

Prof. Fatih Demirci  
Eskişehir, Turkey

Prof. Francesco Epifano  
Chieti Scalo, Italy

Prof. Ana Cristina Figueiredo  
Lisbon, Portugal

Prof. Cristina Gracia-Viguera  
Murcia, Spain

Dr. Christopher Gray  
Saint John, NB, Canada

Prof. Dominique Guillaume  
Reims, France

Prof. Duvvuru Gunasekar  
Tirupati, India

Prof. Hisahiro Hagiwara  
Niigata, Japan

Prof. Judith Hohmann  
Szeged, Hungary

Prof. Tsukasa Iwashina  
Tsukuba, Japan

Prof. Leopold Jirovetz  
Vienna, Austria

Prof. Phan Van Kiem  
Hanoi, Vietnam

Prof. Niel A. Koorbanally  
Durban, South Africa

Prof. Chiaki Kuroda  
Tokyo, Japan

Prof. Hartmut Laatsch  
Gottingen, Germany

Prof. Marie Lacaillle-Dubois  
Dijon, France

Prof. Shoei-Sheng Lee  
Taipei, Taiwan

Prof. M. Soledade C. Pedras  
Saskatoon, Canada

Prof. Luc Pieters  
Antwerp, Belgium

Prof. Peter Proksch  
Düsseldorf, Germany

Prof. Phila Raharivelomanana  
Tahiti, French Polynesia

Prof. Stefano Serra  
Milano, Italy

Dr. Bikram Singh  
Palampur, India

Prof. Leandros A. Skaltsounis  
Zografou, Greece

Prof. John L. Sorensen  
Manitoba, Canada

Prof. Johannes van Staden  
Scottsville, South Africa

Prof. Valentin Stonik  
Vladivostok, Russia

Prof. Winston F. Tinto  
Barbados, West Indies

Prof. Sylvia Urban  
Melbourne, Australia

Prof. Karen Valant-Vetschera  
Vienna, Austria

**INFORMATION FOR AUTHORS**

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

**To Subscribe:** Natural Product Communications is a journal published monthly. 2017 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

## Antiviral Activity of *Aspalathus linearis* against Human Influenza Virus

Ratika Rahmasari<sup>a</sup>, Takahiro Haruyama<sup>a,b</sup>, Siriwan Charyasriwong<sup>a</sup>, Tomoki Nishida<sup>a</sup> and Nobuyuki Kobayashi<sup>a,b\*</sup>.

<sup>a</sup>Laboratory of Molecular Biology of Infectious Agents, Graduate School of Biomedical Sciences, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan

<sup>b</sup>Central Research Center, AVSS Corporation, 1-22 Wakaba-machi, Nagasaki 852-8137, Japan.

nobnob@nagasaki-u.ac.jp

Received: January 10<sup>th</sup>, 2017; Accepted: February 16<sup>th</sup>, 2017

Influenza A viruses are responsible for annual epidemics and occasional pandemics, which cause significant morbidity and mortality. The limited protection offered by influenza vaccination, and the emergence of drug-resistant influenza strains, highlight the urgent need for the development of novel anti-influenza drugs. However, the search for antiviral substances from the library of low molecular weight chemical compounds is limited. Thus, because of their natural diversity and accessibility, plants or plant-derived materials are rapidly becoming valuable sources for the discovery and development of new antiviral drugs. In this study, crude extracts of *Aspalathus linearis*, a plant reported to have anti-HIV activity, were evaluated *in vitro* for their activity against the influenza A virus. Of the extracts tested, an alkaline extract of *Aspalathus linearis* demonstrated the strongest inhibition against influenza A virus and could also inhibit different types of influenza viruses, including Oseltamivir-resistant influenza viruses A and B. Our time course of addition studies indicated that the alkaline extract of *Aspalathus linearis* exerts its antiviral effect predominantly during the late stages of the influenza virus replication process.

**Keywords:** *Aspalathus linearis*, Alkaline extract, Anti-influenza activity, Natural product.


Influenza viruses are enveloped, single-stranded RNA viruses, belonging to the *Orthomyxoviridae* family. The genetic reassortment of influenza virus has resulted in the emergence of novel antigenic subtypes of the influenza virus such as 2009 H1N1 swine and 2013 H7N9 avian influenza virus, which can cause infection in humans [1a-c]. To prevent infection, annual vaccination is highly recommended. However, the relative benefits of the “flu” vaccine are significantly reduced when the antigenic component of this vaccine is miss-matched with the circulating virus [1a-e]. Consequently, the widespread use of an appropriate anti-influenza virus drug is essential to control the spread of the influenza virus. Currently, two main classes of anti-influenza drug are available for the prevention and treatment of influenza virus; the neuraminidase (NA) inhibitor and the matrix 2 protein (M2) ion channel inhibitor. However, because of the high mutation rates of influenza genomes, reports already describe the rapid and widespread emergence of resistance to the M2 ion channel inhibitor; furthermore, resistance to the NA inhibitor has also been reported for seasonal influenza virus A [for review, see 2a-b]. The emergence of such drug resistance is a major driving factor for the development of novel anti-influenza drugs.

However, the discovery of new drugs is facing significant challenges, largely due to a reduction in the number of approved new drugs, and the increasing costs associated with this practice. While low molecular libraries, based on combinatorial chemistry, have provided fresh hope for better success, this has generally failed to improve success rates in terms of the number of new drugs discovered [for review see 3a-b]. Due to their natural diversity and accessibility, plants and plant-derived phytochemicals are rapidly emerging into the limelight in the search for new antiviral [for review see 3c]. Consequently, the antiviral activity of traditional medicines, and their constituents, is becoming of increasing interest for study.

*Aspalathus linearis* is a leguminous plant belonging to the fabacea family which is native to the western-cape of South Africa. Mixtures of dried leaves and flowers from this particular plant, known commonly as Rooibos tea, have been used as traditional medicines and a common health beverage in Europe, Africa, and Japan [for review see 4a]. Traditionally, *A. linearis* is used to cure infant colic, improve appetite, reduce nervous tension and heartburn, and alleviate dermatological problems such as eczema. Its current use as a common health beverage is particularly popular because of the absence of alkaloids, particularly caffeine, and a low concentration of tannin [for review see 4b-c].

*A. linearis* has been reported to contain high amounts of phenols such as aspalathin, nothofagin, isoorientin, orientin, vixetin and rutin, along with a much lower content of quercetin and luteolin [5a-c]. Numerous studies have demonstrated the biological activity of *A. linearis*. Of particular note is that this plant has been reported to exhibit anti-mutagenic activity of aflatoxin B1 and 2-acetylaminofluorene in *Salmonella typhimurium* [6a], anti-cancer activity in rat liver by reducing the number of loci in liver cells [6b], and anti-rotavirus activity [6c] which was attributed to its phenol content. It has also been reported that an aqueous extract of *A. linearis* exhibits antioxidant activity which interferes with stress-reducing activity in rats in which stress was induced by immobilization [6d]. Nakano et al. further showed that the polysaccharide content of the alkaline extract of *A. linearis* could interfere with the binding of HIV-1 to MT-4 cells [7]. In addition, previous studies have reported that quercetin was able to interfere with influenza virus infection [8a-b]. In the present study, considering the emergence of novel antigenic subtypes of the influenza virus and anti-influenza drug resistance, we investigated the anti-influenza virus activity of *A. linearis* extracts *in vitro*.

Phytochemical screening of *A. linearis* against the influenza virus was carried out using extracts of 1% sodium bicarbonate (alkaline)



**Figure 1:** The effect of *Aspalathus linearis* extracts against influenza A virus infection. Evaluation of the activity of each extract against influenza A was performed as described in the experimental section. The extracts were added to MDCK cells in the absence (closed circles) or presence (open circles) of MOI 0.002 A/WSN/33 and incubated at 37°C for 72 h. The relative OD value (%) was also calculated as mentioned earlier. Data are represented as the mean  $\pm$  standard error of mean from two independent experiments (each in triplicate). Extract concentration is represented as % (v/v).

solution, which was prepared as described in a previous publication [7], 70% ethanol (ethanol), ethyl acetate and n-hexane solvent, as described in the experimental section. The cellular toxicity and anti-influenza activity of *A. linearis* extracts were evaluated using the crystal violet (CV) method which represents a quick and simple technique for the quantification of cell death. This method uses Madin-darby canine kidney (MDCK) cells which undergo cell death following infection with the influenza virus; they then lose their adherence and are subsequently detached from the culture dish, thus reducing the amount of crystal violet staining. Inhibition of virus-induced cell death prevents cells from detachment, leading to high levels of crystal violet staining when compared to negative controls.

The antiviral activity of *A. linearis* extracts against influenza virus A/WSN/33 was tested using a multiplicity of infection (MOI) of 0.002. Data showed that alkaline and ethanol extracts could inhibit influenza virus A/WSN/33 infection with an  $IC_{50}$  of  $1.6 \pm 0.2\%$  and  $0.2 \pm 0.1\%$ , respectively (Figure 1). In addition, the second peak observed at 2.5% concentration of ethanol extract (Figure 1B) was caused by cellular toxicity. The cellular toxicity of *A. linearis* extracts was also determined; the  $CC_{50}$  of the alkaline was  $34.6 \pm 8.0\%$  (Supplementary data 1) and ethanol extracts was  $1.3 \pm 0.6\%$ , respectively. However, no toxicity was observed when cells were treated with ethyl acetate and n-hexane extracts.

**Table 1:** Anti-influenza A virus activity of *Aspalathus linearis* extracts.

Extraction	Anti IFV A/WSN/33 activity		
	$IC_{50}$ (%) <sup>a</sup>	$CC_{50}$ (%) <sup>b</sup>	SI <sup>c</sup>
Alkaline	$1.6 \pm 0.2$	$34.6 \pm 8.0$	21.9
Ethanol	$0.2 \pm 0.1$	$1.3 \pm 0.6$	7
Ethyl acetate	-	> 10	-
n-Hexane	-	> 10	-
Oseltamivir	$3.0 \pm 0.0^d$	>1000 <sup>d</sup>	>329

<sup>a</sup> $IC_{50}$  is 50% inhibitory concentration of extract. <sup>b</sup> $CC_{50}$  is 50% cytotoxic concentration of extract. <sup>c</sup>SI is selectivity index which is the ratio of  $CC_{50}$  to  $IC_{50}$ . <sup>d</sup>Oseltamivir unit is  $\mu$ M. All data are represented as mean  $\pm$  standard error of mean of two independent experiments (each in triplicate).

The alkaline extract of *A. linearis*, which possessed a high  $IC_{50}$  value, exhibited less cellular toxicity compared to the ethanol extract, resulting in a higher selectivity index value (SI), (Table 1). In the present study, the alkaline extract of *A. linearis* possessed an  $IC_{50}$  of  $1.6 \pm 0.2\%$  (v/v) with an SI value of 21.9. The alkaline extract of *A. linearis* was duly selected for further evaluation due to its favorable SI value. The alkaline extract of *A. linearis* also exhibited dose dependent activity against A/WSN/33 MOI 0.001 ( $IC_{50}$  =  $1.4 \pm 0.8\%$ ) and MOI 0.1 ( $IC_{50}$  =  $2.7 \pm 1.4\%$ ) infection (data not shown). These results suggest that the alkaline extract of *A. linearis* had potent anti-influenza activity.

We further evaluated the antiviral activity of alkaline extract against different types of human influenza viruses in order to determine its spectrum of activity. For this purpose, a set of laboratory-adapted influenza viruses (A/PR/8/34, B/Brisbane/60/2008, B/Lee/40) and clinical isolates (A/Nagasaki/HA-6/2009, A/Nagasaki/HA-16/2009, A/Nagasaki/HA-33/2009, A/Nagasaki/HA-58/2009 Oseltamivir-resistant, B/Okinawa/4/2014 Oseltamivir-resistant, B/Okinawa/6/2014, B/Okinawa/7/2014 Oseltamivir-resistant, B/Okinawa/8/2014) were examined. These experiments showed that the alkaline extract of *A. linearis* was effective against all of the influenza A and B viruses tested. More importantly, the extract could also inhibit all clinical isolates of the influenza virus including Oseltamivir-resistant virus (Table 2). These results suggested that the *A. linearis* extract had a broad spectrum of antiviral activity.

**Table 2:** Activity of alkaline extract against different types of influenza virus.


Influenza virus	$IC_{50}$ (%)	SI <sup>a</sup>
A/Nagasaki/HA-6/2009	$0.3 \pm 0.0$	134.6
A/Nagasaki/HA-16/2009	$1.3 \pm 0.7$	31.3
A/Nagasaki/HA-33/2009	$1.2 \pm 0.6$	32.4
A/PR/8/34	$0.5 \pm 0.1$	102.9
Oseltamivir-Sensitive		
A/WSN/33	$1.6 \pm 0.2$	21.9
B/Brisbane/60/2008	$3.2 \pm 0.0$	13.5
B/Okinawa/6/2014	$3.0 \pm 1.2$	16.7
B/Okinawa/8/2014	$4.7 \pm 0.8$	12.8
B/Lee/40	$3.1 \pm 0.3$	17.3
Oseltamivir-Resistant		
A/Nagasaki/HA-58/2009	$0.9 \pm 0.0$	43.8
B/Okinawa/4/2014	$3.8 \pm 0.7$	12.4
B/Okinawa/7/2014	$3.2 \pm 0.5$	14.5

<sup>a</sup>SI was calculated as ratio of  $CC_{50}$  from table 1 to each  $IC_{50}$  value. All data are represented as mean  $\pm$  standard error of mean of two independent experiments (each in triplicate).

Oseltamivir-resistant viruses have rapidly emerged across the world for both influenza viruses A and B [for review see 2a-b]. In the current study, the alkaline extract of *A. linearis* showed an inhibitory effect against three clinical isolates which had previously proved resistant to Oseltamivir: A/Nagasaki/HA-58/2009 which harbors the H257Y mutation in the NA glycoprotein [11] and B/Okinawa/4/2009 and B/Okinawa/7/2009 [12b], (Table 2). The ability of the alkaline extract to inhibit Oseltamivir-resistant virus is suggestive evidence that *A. linearis* would be a potential source for development as a novel anti-influenza therapeutic.

To characterize the point within the influenza virus infection cycle which is inhibited by the alkaline extract of *A. linearis*, it was necessary to perform a time course addition study, as reported previously [8a, 12] with some modifications. Five percent of alkaline extract, or 20  $\mu$ M of Oseltamivir, was added into the cells at different time points after virus inoculation at a MOI of 0.001, as depicted in Figure 2. The viral yield from treatment, after infection, was then compared to treatment during infection and Oseltamivir was used as a control drug. This showed that the addition of alkaline extract at 0–12 h could not significantly reduce the relative virus yield when compared to treatment during

infection. In contrast, there was a significant reduction in the yield of virus ( $p < 0.05$ ) when the alkaline extract was added 9 – 12 h after infection (Table 3). As a control, treatment with 20  $\mu$ M of Oseltamivir, which can inhibit NA activity during the later stages of influenza infection, at 0 – 12 h and 9 – 12 h, was shown to significantly reduce the yield of virus. These results indicated that the alkaline extract had the ability to inhibit the influenza virus at late stages of the viral life cycle.



**Figure 2:** Time-of-addition assay used to identify the steps of the influenza A virus life cycle which can be inhibited by alkaline extract. A 5% alkaline extract or 20  $\mu$ M of Oseltamivir was added into infected MDCK cells during infection or at different points of time after infection.

A previous report [7], which examined the antiviral activity of the alkaline extract of *A. linearis* against HIV-1 infection, concluded that the active component of this extract, in this particular instance, was polysaccharide. On the other hand, the active constituent of the alkaline extract of *A. linearis*, which is responsible for inhibiting influenza virus infection, is presently unknown. However, since anti-influenza virus activity could only be observed in ethanol and alkaline extracts (Figure 1), the phenolic content present in the alkaline extract could represent a promising candidate for the active compound [5c, 9a-c]. In addition, Nakano et al. mentioned that the alkaline extract of *A. linearis* contains uronic acid, which was also reported by Mann et al., which is effective in inhibiting influenza sialidase [7, 13]. Our present results suggested that the alkaline extract of *A. linearis*, extracted using the same method as that described by Nakano et al. [7], has potent inhibitory effects against the influenza virus at late stages of infection. Further studies are now needed to determine the active compound(s) involved and their precise mechanism of action.

**Table 3:** Relative virus yield of time of addition assay.

	Relative virus yield (%) <sup>a</sup>	
	5% Alkaline extract	20 $\mu$ M Oseltamivir
During infection	100	100
0 - 12 hpi <sup>b</sup>	28 $\pm$ 17.7	2 $\pm$ 1.6*
0 - 3 hpi	>100	49 $\pm$ 24
3 - 6 hpi	>100	>100
6 - 9 hpi	61 $\pm$ 21	51 $\pm$ 28
9 - 12 hpi	7 $\pm$ 1.4*	1 $\pm$ 0.8*

<sup>a</sup>Relative virus yield was calculated as percentage ratio of treatment after infection to treatment during infection. <sup>b</sup>hpi is abbreviation of hours post infection. \*The asterisks indicate the significant difference between treatment during infection and different time course treatment after infection,  $p < 0.05$ .

## Experimental

**Cells, viruses and samples:** MDCK cells were grown and maintained in Eagle's minimum essential medium (MEM), containing 100 units/ml of penicillin G, 100  $\mu$ g/ml of streptomycin sulfate and supplemented by 5% fetal bovine serum (FBS), at 37°C in 5% CO<sub>2</sub> atmosphere. The influenza A/Wisconsin/33 (A/WSN/33) was then propagated in MDCK cells. The influenza A/Puerto Rico/8/34 (A/PR/8/34) was propagated in MDCK cells in the presence of 2.5  $\mu$ g/ml trypsin (Sigma-Aldrich Co., St. Louis, MO) and the culture supernatants were stored at -80°C. The clinical

isolates of influenza virus from 2009 seasonal patient in Japan (A/Nagasaki/HA-6/2009, A/Nagasaki/HA-16/2009, A/Nagasaki/HA-33/2009, and A/Nagasaki/HA-58/2009 Oseltamivir-resistant) were propagated in a ten day-old embryonic egg for 48 h and then allantoic fluid was collected and stored at -80°C. The laboratory-adapted strains of influenza B/Brisbane/60/2008, B/Lee/40 and clinical isolates of influenza B virus from Japan (B/Okinawa/4/2014 Oseltamivir-resistant, B/Okinawa/6/2014, B/Okinawa/7/2014 Oseltamivir-resistant, and B/Okinawa/8/2014) were also propagated in MDCK cells and culture supernatants stored at -80°C. Viral titer was determined by calculating the 50% tissue culture infectivity dose (TCID<sub>50</sub>). Oseltamivir sulfate was purchased from F. Hoffmann-La Roche Ltd (Basel, Switzerland) and dissolved in ultrapure water to a concentration of 24 mM. Dried *A. linearis* leaves were purchased from All tea RTU Co. Ltd (Nagano, Japan).

**Extract preparation:** The 1% sodium bicarbonate extraction procedure was performed as described previously by Nakano et al. [7] with some modifications. In brief, 1 g of *A. linearis* was extracted with 10 ml of boiled ultrapure water and shaken at 85°C for 3 h. After filtration with a 0.45  $\mu$ m filter, the residue was collected and further extracted in 15 ml of ultrapure water containing 1% sodium bicarbonate at 37°C for 4 h. After re-filtration using a 0.45  $\mu$ m filter, the supernatant (alkaline extract) was finally stored at -80°C.

For ethanol, ethyl acetate and n-hexane extraction, 1 g of *A. linearis* was extracted with 10 ml of 70% ethanol (or ethyl acetate or n-hexane, as appropriate) and shaken for 20 h at room temperature, followed by three separate centrifugation steps at 8740  $\times$ g (Beckman J2-21 centrifuge, JA-20 rotor) for 20 min. The collected supernatant was then evaporated and the pellet diluted with 1% DMSO to become ethanol (or ethyl acetate or n-hexane) extract. All extracts were stored at -80°C to await further analysis.

## Cellular toxicity and anti-influenza activity of *A. linearis* extracts:

Cellular toxicity was evaluated using a 96 well-culture plate. Confluent MDCK cells were first seeded and incubated overnight. The cell supernatant was then aspirated and washed by serum-free MEM, followed by the addition of 100  $\mu$ l of serum-free MEM containing 1% vitamins and 100  $\mu$ l of serial two-fold dilution of extracts. The mixtures were then incubated at 37°C in an atmosphere containing 5% CO<sub>2</sub> for 3 days. The supernatant was then aspirated and the cells fixed with 70% ethanol and then stained with 0.5% CV solution. Cells were then dried and absorbance measured using an Infinite M200 Tecan plate reader. The CC<sub>50</sub> was then calculated by linear regression analysis using Microsoft Excel software.

The anti-influenza assay was carried out using a 96 well-culture plate. Confluent MDCK cells were seeded and incubated at 37°C overnight. The supernatant was then aspirated and washed with serum-free MEM, followed by the addition of 100  $\mu$ l serial dilutions of extracts and 100  $\mu$ l of influenza virus A (A/PR/8/34 or A/WSN/33 or A/Nagasaki/HA-6/2009 or A/Nagasaki/HA-16/2009 or A/Nagasaki/HA-33/2009 or A/Nagasaki/HA-58/2009 Oseltamivir-resistant) or influenza virus B (B/Brisbane/60/2008 or B/Lee/40 or B/Okinawa/4/2014 Oseltamivir-resistant or B/Okinawa/6/2014 or B/Okinawa/7/2014 Oseltamivir-resistant or B/Okinawa/8/2014) solution equivalent to a MOI of 0.002. The mixtures were then incubated at 37°C in an atmosphere containing 5% CO<sub>2</sub> for 3 days. The supernatant was then aspirated and the cells fixed with 70% ethanol and stained with 0.5% CV. Cells were dried and absorbance measured as described above. The relative optical

density (OD) value was calculated from the ratio of absorbance in the presence of virus to absorbance in the absence of virus. The  $IC_{50}$  was then calculated by linear regression analysis using Microsoft Excel software. The SI was then determined by the ratio of  $CC_{50}$  to  $IC_{50}$  values.

**The time-of-addition assay:** MDCK cells were seeded at a concentration of  $2 \times 10^5$  cells per well into 24 well-culture plates and incubated overnight. The cell monolayer was then infected with A/WSN/33 at MOI 0.001 and treated with a 5% alkaline extract of *A. linearis*, or 20  $\mu$ M of Oseltamivir at different times of addition as follows: a) treatment during infection: 100  $\mu$ l of 5% alkaline extract of *A. linearis* or 20  $\mu$ M of Oseltamivir and virus equal to MOI 0.001 were added to the cells simultaneously and incubated at 37°C for 1 h. Unattached viruses were removed and the cells washed twice with MEM followed by the addition of 1 ml of MEM; b) treatment after infection: cells were infected by 100  $\mu$ l of virus

equal to MOI 0.001 at 37°C for 1 h. Unattached viruses were removed and cells washed twice with MEM. Following infection, 1 ml of MEM containing 5% alkaline extract of *A. linearis*, or 20  $\mu$ M of Oseltamivir, were added at different time periods, as shown in Figure 2. Twelve hours after infection, the supernatant from different time points of alkaline extract or Oseltamivir treatment were collected and subjected to viral titration using the 50 % tissue culture infectious dose assay. The viral titer was then calculated using the Reed-Muench method. The relative viral yield value was calculated as the percentage ratio of treatment after infection relative to treatment during infection.

**Statistical analysis:** Data are represented as the mean  $\pm$  standard error of the mean from two independent experiments (each in triplicate). The comparison of test samples and controls was performed using the student's *t* test. A *p* value < 0.05 was considered as statistically significant.

## References

- [1] (a) <http://www.cdc.gov/flu/professionals/vaccination/effectivenessqa.htm>; (b) [http://www.who.int/influenza/human\\_animal\\_interface/influenza\\_h7n9/en/](http://www.who.int/influenza/human_animal_interface/influenza_h7n9/en/); (c) [https://www.cdc.gov/h1n1flu/information\\_h1n1\\_virus\\_qa.htm](https://www.cdc.gov/h1n1flu/information_h1n1_virus_qa.htm); (d) Pica N, Palese P. (2013) Toward a universal influenza virus vaccine: Prospects and challenges. *Annual Review of Medicine*, **64**, 189–202; (e) <http://www.cdc.gov/flu/about/viruses/change.htm>
- [2] (a) Frederick G, Hayden FG, and De Jong MD. (2011) Emerging influenza antiviral resistance threats. *The Journal of Infectious Diseases*, **203**, 6–10; (b) McKimm-Breschkin. (2012) Influenza neuraminidase inhibitors: Antiviral action and mechanisms of resistance. *Influenza and Other Respiratory Viruses*, **7**, 25–36.
- [3] (a) Katiyar C, Gupta A, Kanjilal S, Katiyar S. (2012) Drug discovery from plant sources: An integrated approach. *Ayu*, **33**, 10-33; (b) Atanasov AG, Waltenberger B, Pferschy-wenzig EM, Linder T, Wawrosch C, Uhrin P, Temml V, Wang L, Schwaiger S, Heiss EH, Rollinger JM, Schuster D, Breuss JM, Bochkov V, Mihovilovic MD, Kopp B, Bauer R, Dirsch VM and Stuppner H. (2015) Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology Advances*, **33**, 1582-1614; (c) Kurokawa M, Shimizu T, Watanabe W, Shiraki K. (2010) Development of new antiviral agents from natural product. *The Open Antimicrobial Agents Journal*, **2**, 49-57.
- [4] (a) Mackay DL, Blumberg JB. (2007) Review of bioactivity of South African herbal tea: Rooibos tea (*Aspalathus linearis*) and Honey bush (*Cyclopia intermedia*). *Phytotherapy Research*, **21**, 1-16; (b) Joubert E, Gelderblom WCA, Louw A, de Beer D. (2008) South African herbal teas: *Aspalathus linearis*, *Cyclopia* spp. and *Athrixia phylicoides*-A review. *Journal of Ethnopharmacology*, **119**, 376–412; (c) Morton JF. (1983) Rooibos Tea, *Aspalathus linearis*, a caffeine-less, low-tannin beverage. *Economic Botany*, **37**, 64-173.
- [5] (a) Joubert E. (1995) HPLC quantification of the dihydrochalcones, aspalathin and nothofagin in Rooibos tea (*Aspalathus linearis*) as affected by processing. *Food Chemistry*, **55**, 403-411; (b) Bramati L, Aquilano F, Pietta P. (2003) Unfermented Rooibos Tea: Quantitative characterization of flavonoids by HPLC-UV and determination of the total antioxidant activity. *Journal of Agricultural and Food Chemistry*, **51**, 7472–7474; (c) Joubert E, de Beer D. (2012) Phenolic content and antioxidant activity of rooibos food ingredient extracts. *Journal of Food Composition and Analysis*, **27**, 45-51; (e) KrafczykN, Glomb MA. (2008) Characterization of phenolic compounds in Rooibos tea. *Journal of Agricultural and Food Chemistry*, **56**, 3368-3376.
- [6] (a) Van Der Merwe JD, Joubert E, Richards ES, Manley M, Snijman PW, Marnewick JL, Gelderblom WC. (2006) A comparative study on the antimutagenic properties of aqueous extracts of *Aspalathus linearis* (rooibos), different *Cyclopia* spp. (honeybush) and *Camellia sinensis* teas. *Mutation Research*, **611**, 42–53; (b) Marnewick JL, Van Der Westhuizen FH, Joubert E, Swanevelder S, Swart P, Gelderblom WC. (2009) Chemoprotective properties of Rooibos (*Aspalathus linearis*), Honeybush (*Cyclopia intermedia*) herbal and green and black (*Camellia sinensis*) teas against cancer promotion induced by fumonisin B1 in rat liver. *Food and Chemical Toxicology*, **47**, 220–229; (c) Knipping K, Garssen J, Van't LB. (2012) An evaluation of the inhibitory effects against rotavirus infection of edible plant extracts. *Virology Journal*, **9**, 137; (d) Hong I-S, Lee H-Y, Kim H-P. (2014) Anti-oxidative effects of Rooibos tea (*Aspalathus linearis*) on immobilization-induced oxidative stress in rat brain. *PLoS One*, **9**(1), e87061.
- [7] Nakano M, Itoh Y, Mizuno T, Nakashima H. (1997) Polysaccharide from *Aspalathus linearis* with strong Anti-HIV activity. *Bioscience, Biotechnology, and Biochemistry*, **61**, 267–271.
- [8] (a) Wu W, Li R, Li X, He J, Jiang S, Liu S, Yang J. (2015) Quercetin as an antiviral agent inhibits influenza A virus (IAV) entry. *Viruses*, **8**(6), doi:10.3390/v8010006; (b) Choi HJ, Song JH, Kwon DH. (2012) Quercetin 3-rhamnoside exerts anti-influenza A virus activity in mice. *Phytotherapy Research*, **26**, 462-464.
- [9] (a) Nencioni L, Iuvara A, Aquilano K, Ciriolo MR, Cozzolino F, Rotilio G, Garaci E, Palamara AT. (2003) Influenza A virus replication is dependent on an antioxidant pathway that involves GSH and Bcl-2. *FASEB Journal*, **17**, 758-760; (b) Ciriolo MR, Palamara AT, Incerci S, Lafavia E, Buè MC, De Vito P, Garaci E, Rotilio G. (1997) Loss of GSH, oxidative stress, and decrease of intracellular pH as sequential steps in viral infection. *The Journal of Biological Chemistry*, **272**, 2700–2708. (c) Marnewick JL, Joubert E, Swart P, Van Der Westhuizen F, Gelderblom WC. (2003) Modulation of hepatic drug metabolizing enzymes and oxidative status by rooibos (*Aspalathus linearis*) and honey bush (*Cyclopia intermedia*), Green and Black (*Camellia sinensis*) teas in rats. *Journal of Agricultural and Food Chemistry*, **51**, 8113-8119.
- [10] (a) Sakagami H, Ohkoshi E, Amano S, Satoh K, Maki J, Kanamoto T, Terakubo S, Nakashima H, Sunaga K, Otsuki T, Ikeda H and Fukuda T. (2013) Efficient utilization of plant resources by alkaline extraction. *Alternative and Integrative Medicine*, **2**(133), doi:10.4172/2327-5162.1000133; (b) Huang SQ, Li JW, Wang Z, Pan HX, Chen JX, Ning ZX. (2010) Optimization of alkaline extraction of polysaccharide from *Ganoderma lucidum* and their effect on immune function in mice. *Molecules*, **15**, 3694-3708.
- [11] Kawano H, Haruyama T, Hayashi Y, Sinoda M, Kobayashi N. (2011) Genetic analysis and phylogenetic characterization of pandemic (H1N1) 2009 influenza viruses that found in Nagasaki, Japan. *Japanese Journal of Infectious Diseases*, **64**, 195-203.
- [12] (a) Makau JN, Watanabe K, Kobayashi N. (2013) Anti-influenza activity of *Alchemilla mollis* extract: Possible virucidal activity against influenza virus particles. *Drug Discoveries & Therapeutics*, **7**, 189-195; (b) Charyasriwong S, Haruyama T, Kobayashi N. (2016) *In vitro* evaluation of the antiviral activity of methylglyoxal against influenza B virus infection. *Drug Discoveries and Therapeutics*, **10**, 201-210.
- [13] Mann MC, Islam T, Dyason JC, Florio P, Trower CJ, Thompson RJ, Itzstein MV. (2006) Unsaturated *N*-Acetyl-D-glucosaminuronic acid glycosides as inhibitors of influenza virus sialidase. *Glycoconjugate Journal*, **23**, 127-133.

<b>GC-MS-based Metabolomic Profiling of Thymoquinone in Streptozotocin-induced Diabetic Nephropathy in Rats</b> Mohammad Raish, Ajaz Ahmad, Basit L. Jan, Khalid M. Alkharfy, Kazi Mohsin, Syed Rizwan Ahamad and Mushtaq Ahmad Ansari	553
<b>Synthesis and Bioconversion of Curcumin Analogs</b> Glenroy D. A. Martin, Cameron McKenzie and Monica Moore	559
<b>Chemical Constituents of <i>Chirita drakei</i></b> Nguyen Thi Hoang Anh, Nguyen Van Tuan, Dao Duc Thien, Tran Duc Quan, Nguyen Thanh Tam, Giang Thi Kim Lien, Katrin Franke, Trinh Thi Thuy and Tran Van Sung	563
<b>Antibacterial Activity of 2-(3',5'-Dibromo-2'-methoxyphenoxy)-3,5-dibromophenol Isolated from <i>Phyllospongia papyracea</i></b> Shi Sun, Corene Canning, Kuiwu Wang, Wenjun Zhu, Fei Yang, Yifan Zhang and Kequan Zhou	567
<b>The Antitumor Antibiotics Complex of Aureolic Acids from the Marine Sediment-associated Strain of <i>Streptomyces</i> sp. KMM 9048</b> Nataliya I. Kalinovskaya, Lyudmila A. Romanenko, Anatoly I. Kalinovskiy, Svetlana P. Ermakova, Pavel S. Dmitrenok and Shamil Sh. Afiyatulloev	571
<b>Anti-dormant Mycobacterial Activity of Viomellein and Xanthomegnin, Naphthoquinone Dimers Produced by Marine-derived <i>Aspergillus</i> sp.</b> Kentaro Kamiya, Masayoshi Arai, Andi Setiawan and Motomasa Kobayashi	579
<b>Two New Sulfonoglycolipids from the Green Alga <i>Codium dwarkense</i></b> Liaqat Ali, Lubna Al-Kharusi and Ahmed Al-Harrasi	583
<b>A Convergent Total Synthesis of Balticolid</b> Avuluri Srilatha, Jhillu S. Yadav and Basi V. Subba Reddy	587
<b>Improvement Effects of Wasabi (<i>Wasabia japonica</i>) Leaves and Allyl Isothiocyanate on Stomach Lesions of Mongolian Gerbils Infected with <i>Helicobacter pylori</i></b> Shuichi Masuda, Hideki Masuda, Yuko Shimamura, Chitose Sugiyama and Fumiyo Takabayashi	595
<b>Antiviral Activity of <i>Aspalathus linearis</i> against Human Influenza Virus</b> Ratika Rahmasari, Takahiro Haruyama, Siriwan Charyasriwong, Tomoki Nishida and Nobuyuki Kobayashi	599
<b>Methyl 3-(5-(prop-1-yn-1-yl)thiophen-2-yl)propanoate: A Rare Acetylene Derivative from <i>Artemisia absinthium</i> Root Essential Oil</b> Polina D. Blagojević, Marko S. Pešić and Niko S. Radulović	603
<b>Chemical Investigation on Leaf, Flower and Fruit Oils of <i>Lantana camara</i> from Côte d'Ivoire</b> Fatimata Nea, Evelyne Amenan Tanoh, Thierry Acafou Yapi, Gabriel Garcia, Felix Tomi and Zanahi Félix Tonzibo	607
<b>Chemical Composition and Anti-inflammatory Activity of Algerian <i>Thymus vulgaris</i> Essential Oil</b> Wafae Abdelli, Fouad Bahri, Abderrahmane Romane, Martina Höferl, Juergen Wanner, Erich Schmidt and Leopold Jirovetz	611
<b><i>Elionurus tristis</i> Essential Oil: GC-MS Analysis and Antioxidant and Antituberculosis Activities</b> Brice Hervé Yedomon, Isabelle Saves, Narjes Mtimet, Emmanuel Guy Raelison, Patricia Constant, Mamadou Daffé and Jalloul Bouajila	615
<b>Chemical Composition and Bactericidal Activity of the Essential Oils of Four Species of Annonaceae Growing in Brazilian Amazon</b> Joelma Moreira Alcântara, Juliana Mesquita V. M. de Lucena, Roselaine Facanali, Marcia Ortiz M. Marques and Maria da Paz Lima	619
<b>Coriander (<i>Coriandrum sativum</i>) Essential Oil: Effect on Multidrug Resistant Uropathogenic <i>Escherichia coli</i></b> Francesca Scazzocchio, Lorenzo Mondì, Maria Grazia Ammendolia, Paola Goldoni, Antonella Comanducci, Massimiliano Marazzato, Maria Pia Conte, Federica Rinaldi, Maria Elisa Crestoni, Caterina Frascchetti and Catia Longhi	623

### Accounts/Reviews

<b>Dietary Soy Isoflavone: A Mechanistic Insight</b> Renu Chadha, Yashika Bhalla, Ankita Jain, Kunal Chadha and Maninder Karan	627
<b>Distribution and Diversity of Usage of the Amaryllidaceae in the Traditional Remediation of Infectious Diseases</b> Jerald J. Nair, Johannes van Staden, Susanna L. Bonnet and Anke Wilhelm	635



# Natural Product Communications

## 2017

Volume 12, Number 4

### Contents

<u>Original Paper</u>	<u>Page</u>
<b>Spasmolytic Action of Preparations and Compounds from <i>Hofmeisteria schaffneri</i></b> Araceli Pérez-Vásquez, Guadalupe Ángeles-López, Isabel Rivero-Cruz, Laura Flores-Bocanegra, Edelmira Linares, Robert Bye and Rachel Mata	475
<b>Sesquiterpene Quinones and Diterpenes from <i>Smenospongia cerebriformis</i> and Their Cytotoxic Activity</b> Le Thi Huyen, Dan Thuy Hang, Nguyen Xuan Nhiem, Bui Huu Tai, Hoang Le Tuan Anh, Tran Hong Quang, Pham Hai Yen, Chau Van Minh, Nguyen Van Dau and Phan Van Kiem	477
<b>Comparison of Different Strains of <i>Agrobacterium rhizogenes</i> for Hairy Root Induction and Betulin and Betulinic Acid Production in <i>Morus alba</i></b> Chang Ha Park, Shicheng Zhao, Hyeon Ji Yeo, Ye Eun Park, Thanislas Bastin Baskar, Mariadhas Valan Arasu, Naif Abdullah Al-Dhabi and Sang Un Park	479
<b>Two New Triterpenes and Other Compounds from <i>Mentha aquatica</i> (Lamiaceae)</b> Maria Ferhat, Zahia Kabouche, Yoshinori Fujimoto and Hiroshi Araya	483
<b>Chemical Composition and Antioxidant Activity of <i>Lawsonia inermis</i> Seed Extracts from Morocco</b> Issmail Nounah, Ahmed Hajib, Hicham Harhar, Nadia El Madani, Saïd Gharby, Dom Guillaume and Zoubida Charrouf	487
<b>Talarofuranone, a New Talaroconvolutin Analog from the Endophytic Fungus <i>Talaromyces purpurogenus</i> from <i>Pouteria campechiana</i> Seeds</b> K.G.E. Padmathilake, H.M.S.K.H. Bandara, M. Mallique Qader, N. Savitri Kumar, Lalith Jayasinghe, Hironori Masubuti and Yoshinori Fujimoto	489
<b>Ochroborbone, A New Cytotoxic Indole Alkaloid from <i>Ochrosia borbonica</i></b> Yan-Ping Liu, A-Hong Chen, Ruo-Heng Li, Hui-Wen Yang, Hai-Nan Bao, Liang Lai, Kang Zong and Yan-Hui Fu	491
<b>A New Ajmaline-type Alkaloid from the Roots of <i>Rauvolfia serpentina</i></b> Thitima Rukachaisirikul, Suwadee Chokchaisiri, Parichat Suebsakwong, Apichart Suksamrarn and Chainarong Tocharus	495
<b>Positions of Hydroxyl Groups in Chrysin are Critical for Inhibiting Plasminogen Activator Inhibitor-1 Release from Human Umbilical Vein Endothelial Cells</b> Naoki Ohkura, Kumiko Ando, Yuko Takata, Shiho Kanai, Kenichi Ishibashi, Masahiko Taniguchi, Tomoki Tatefuji and Gen-ichi Atsumi	499
<b>Constituents of <i>Flourensia blakeana</i> (Asteraceae)</b> Mila Arjona, Alejandro A. Alarcón, Adriana Pacciaroni and Rosana Alarcón	503
<b>Monoamine Oxidase Inhibitory Activity of Biflavonoids from Branches of <i>Garcinia gardneriana</i> (Clusiaceae)</b> Maria Angélica Recalde-Gil, Luiz Carlos Klein-Júnior, Carolina dos Santos Passos, Juliana Salton, Sérgio Augusto de Loreto Bordignon, Franco Delle Monace, Valdir Cechinel Filho and Amélia Teresinha Henriques	505
<b>Effects of Intestinal Microecology on Metabolism and Pharmacokinetics of Oral Wogonoside and Baicalin</b> Shihua Xing, Mengyue Wang, Ying Peng and Xiaobo Li	509
<b>Protoflavanones from the Wood Stem of <i>Salvertia convallariodora</i></b> Mariana L. de Mesquita, José E. de Paula, Laila S. Espindola, Luiz A. L. Soares, Tania M. G. da Silva, Celso A. Camara and Telma M. G. Da Silva	515
<b>Chemical Composition and Antioxidant Activity of <i>Cirsium vulgare</i> Inflorescences</b> Jolanta Nazaruk, Sebastian Chłędzik, Jakub Strawa, Katarzyna Bazydło and Anna Wajs-Bonikowska	519
<b>Glycosylation and Methylation of Quercetin and Myricetin by Cultured Cells of <i>Phytolacca americana</i></b> Yuya Fujitaka, Kei Shimoda, Naoji Kubota, Minami Araki, Tatsuya Onishi, Noriyuki Nakayama, Kohji Ishihara, Masato Tanigawa, Hatsuyuki Hamada and Hiroki Hamada	523
<b>Buxusoside, a Flavonoid Disaccharide from <i>Buxus sinica</i></b> Min Wang, Guo-Bo Xu, Jun Liu, Yue-Hao Zhang, Jun-Hong Liu, Jing Li, Meng Zhou, Yao-Hua Xiao, Xun He and Shang-Gao Liao	525
<b>Isoflavone Composition, Total Phenolic Content and Antioxidant Capacity of Soybeans with Colored Seed Coat</b> Mira Bursać, Milica Atanacković Krstonošić, Jegor Miladinović, Đorđe Malenčić, Ljiljana Gvozdenović and Jelena Hogervorst Cvejić	527
<b>Main Constituents and Antidiabetic Properties of <i>Otholobium mexicanum</i></b> Alirica I. Suárez, Zaw Min Thu, Jorge Ramírez, Diana León, Luis Cartuche, Chabaco Armijos and Geovanni Vidari	533
<b>Syntheses of Benzo[c]Chromen-6-ones by Palladium Catalyzed C–H Bond Activation using Diazonium Salts</b> Liang-Zhu Huang, Rui-Yue Ma, Lu-Nan Zhou, Zhen-Ting Du and Tao Zhang	537
<b>Antioxidant Activity and Phenolic Composition of a Red Bean (<i>Phaseolus vulgaris</i>) Extract and its Fractions</b> Ryszard Amarowicz, Magdalena Karamać, Montserrat Dueñas and Ronald B. Pegg	541
<b>Variation in Phenolic Composition of <i>Knautia arvensis</i> in Correlation with Geographic Area and Plant Organ</b> Erna Karalija, Edina Muratović, Petr Tarkowski and Sanja Čavar Zeljković	545
<b>Arbutin Content and Tyrosinase Activity of <i>Bergenia</i> Extracts</b> Lenka Tůmová, Iva Dolečková, Helena Hendrychová and Marie Kašparová	549

Continued inside backcover